Journal of Pharmaceutical and Biomedical Analysis

Volume 12 1994

LIST OF CONTENTS

Number 1

	iii	Editorial
	v	APQ Newsletter
Analytical Survey		
D. T. Burns	1	Cyclodextrins as versatile chiral recognition reagents for use in a variety of optical and separative analyses
Full Papers		
M. J. Lynch, J. Masters, J. P. Pryor, J. C. Lindon, M. Spraul, P. J. D. Foxall and J. K. Nicholson	5	Ultra high field NMR spectroscopic studies on human seminal fluid, seminal vesicle and prostatic secretions
M. G. Donato, W. Baeyens, W. van den Bossche and P. Sandra	21	The determination of non-steroidal antiinflammatory drugs in pharmaceuticals by capillary zone electrophoresis and micellar electrokinetic capillary chromatography
A. F. Casy	27	Rapid identification of ergot derivatives by ^{1}H-NMR spectroscopy
A. F. Casy	41	Mass spectrometry as an aid to the identification of ergots and dihydroergots: comparison of hard and soft ionization techniques
P. Ballard, M. D. Malone and B. Law	47	The use of competitive displacement agents to resolve albumin binding problems observed during the development of a radioimmunoassay for ICI 215001
K. Ensing, I. J. Bosman, A. C. G. Egberts, J. P. Franke and R. A. de Zeeuw	53	Application of radioreceptor assays for systematic toxicological analysis—1. Procedures for radioreceptor assays for antihistaminics, anticholinergics and benzodiazepines
K. Ensing, I. J. Bosman, A. C. G. Egberts, J. P. Franke and R. A. de Zeeuw	59	Application of radioreceptor assays for systematic toxicological analysis—2. Theoretical considerations and evaluation
T. J. Siahaan and K. Lutz	65	Conformational study of cyclo[Gln-Trp-Phe-Gly-Leu-Met] as NK-2 antagonist by NMR and molecular dynamics
I. I. Hewala	73	Gas–liquid chromatographic and difference spectrophotometric techniques for the determination of benzaldehyde in benzyl alcohol
J. Prat, M. Pujol, V. Girona, M. Muñoz and L.-A. Solé	81	Stability of carboplatin in 5% glucose solution in glass, polyethylene and polypropylene containers
B. R. Thomas, X. G. Fang, P. Shen and S. Ghodbane	85	Mixed ion pair liquid chromatography method for the simultaneous assay of ascorbic acid, caffeine, chlorpheniramine maleate, dextromethorphan HBr monohydrate and paracetamol in Frenadol™ sachets
A. M. Di Pietra, R. Gotti, D. Bonazzi, V. Andrisano and V. Cavrini	91	HPLC determination of glutathione and L-cysteine in pharmaceuticals after derivatization with ethacrynic acid
M. A. Faouzi, T. Dine, M. Luyckx, F. Goudaliez, M. L. Mallevais, B. Gressier, C. Brunet, M. Cazin and J. C. Cazin	99	Stability and compatibility studies of cephamandole nafate with PVC infusion bags
K. Wang, R. W. Blain and A. J. Szuna	105	Multidimensional narrow bore liquid chromatography analysis of Ro 24-0238 in human plasma
J.-F. Chaulet, G. Grelaud, P. Bellemin-Magninot, C. Mounier and J.-L. Brazier	111	Simultaneous determination of chloroquine, proguanil and their metabolites in human biological fluids by high-performance liquid chromatography
P. Thongnopnua and K. Viwatwongsa	119	Quantitative analysis of nifedipine in plasma by high-performance liquid chromatography

Short Communications

M. Bodiroga, D. Agbaba, D. Zivanov-Stakic and R. Popovic 127 Determination of pralidoxime chloride in pharmaceutical dosage forms by isotachophoresis

S. Kumbhat 131 Study of ocular penetration of flurbiprofen in human aqueous humour

D. R. Rutledge, A. H. Abadi and L. M. Lopez 135 Liquid chromatographic determination of celiprolol, diltiazem, desmethyldiltiazem and deacetyldiltiazem in plasma using a short alkyl chain silanol deactivated column

P. Nebinger and M. Koel 141 Determination of serum zidovudine by ultrafiltration and high-performance liquid chromatography

Number 2

Full Papers

J.-R. Lin, N. S. Brown, R. A. Newman and D. H. Ho 145 Determination of peplomycin in mouse tissues and biofluids by radioimmunoassay

I. Panderi and M. Parissi-Poulou 151 Simultaneous determination of clopamide–pindolol combination in tablets by zero-crossing derivative spectrophotometry

D.-S. Lee, H.-J. Han, K. Kim, W.-B. Park, J.-K. Cho and J.-H. Kim 157 Dissociation and complexation of fluoroquinolone analogues

R. Andersin, J. Ovaskainen and S. Kaltia 165 Photochemical decomposition of midazolam. III—Isolation and identification of products in aqueous solutions

R. Lozano, J. M. Joseph and B. J. Kline 173 Temperature, pH and agitation rate as dissolution test discriminators of zofenopril calcium tablets

M. Eisman, M. Gallego and M. Varcárcel 179 Determination of papaverine and cocaine by use of a precipitation system coupled on-line to an atomic absorption spectrometer

M. S. Boukraa, D. Deruaz, A. Bannier, M. Desage and J. L. Brazier 185 Detection of ^{13}C labelled compounds by gas chromatography coupled to atomic emission detection—application to caffeine metabolites

C. M. Grosse, I. M. Davis, R. F. Arrendale, J. Jersey and J. Amin 195 Determination of remifentanil in human blood by liquid–liquid extraction and capillary GC–HRMS–SIM using a deuterated internal standard

A. N. Abdelrahman, E. I. A. Karim and K. E. E. Ibrahim 205 Determination of chloroquine and its decomposition products in various brands of different dosage forms by liquid chromatography

S. K. Yang 209 Base-catalysed rearrangement of temazepam

X.-Z. Qin, D. P. Ip, K. H.-C. Chang, P. M. Dradransky, M. A. Brooks and T. Sakuma 221 Pharmaceutical application of LC–MS. 1—Characterization of a famotidine degradate in a package screening study by LC–APCI MS

G. Lhoëst, N. Maton, A. Laurent and R. K. Verbeeck 235 Isolation and identification of a FK-506 C_{36}–C_{37} dihydrodiol from erythromycin-induced rabbit liver microsomes

K. Selinger, C. Lanzo and A. Sekut 243 Determination of remifentanil in human and dog blood by HPLC with UV detection

C. Kohl, H. Oelschläger and D. Rothley 249 Identification of drug glucuronides in human urine by RP-HPLC after derivatization

D. M. Morris and K. Selinger 255 Determination of 2′-deoxy-3′-thiacytidine (3TC) in human urine by liquid chromatography: direct injection with column switching

H. H. Farrish, P.-H. Hsyu, J. F. Pritchard, K. R. Brouwer and J. Jarrett 265 Validation of a liquid chromatography post-column derivatization assay for the determination of cisplatin in plasma

Short Communications

P. Gratteri, S. Furlanetto, S. Pinzauti, E. La Porta, P. Mura and G. Santoni 273 Adsorptive stripping voltammetry for thiomersal assay

A. Sumathi and V. U. Rao 277 Comparison of accelerated stability data for 5-fluorouracil obtained by a newly developed colorimetric method and by the standard spectrophotometric method

M. J. Fossler, J. Leslie, M. Burke, J. Devane and D. Young 281 A reversed-phase ion-pair liquid chromatography method for the determination of 4-aminopyridine in Sprague–Dawley rats

Number 3

Review

T. R. Krishnan and I. Ibraham 287 Solid-phase extraction technique for the analysis of biological samples

Full Papers

J. W. Jaroszewski and A. Olsson 295 Determination of enantiomeric purity of nicotine in pharmaceutical preparations by ^{13}C-NMR in the presence of a chiral lanthanide shift reagent

R. M. E. Richards and D. K. L. Xing 301 Capillary zone electrophoresis assay of the uridine diphosphate *N*-acetylmuramyl peptide precursors and the disaccharide pentapeptide derivative of bacterial cell wall peptidoglycan

A. Memoli, L. G. Palermiti, V. Travagli and F. Alhaique 307 Effects of surfactants on the spectral behaviour of calcein

J. L. Vilchez, G. Sánchez-Palencia, R. Avidad and A. Navalón 313 Determination of boldine in drug formulations by first-derivative synchronous spectrofluorimetry

P. L. Weber, T. J. O'Shea and S. M. Lunte 319 Separation and quantitation of the amino acid neurotransmitters in rat brain by capillary electrophoresis

H. P. Hendrickson, A. D. Kaufman and C. E. Lunte 325 Electrochemistry of catechol-containing flavonoids

H. P. Hendrickson, M. Sahafayen, M. A. Bell, A. D. Kaufman, M. E. Hadwiger and C. E. Lunte 335 Relationship of flavonoid oxidation potential and effect on rat hepatic microsomal metabolism of benzene and phenol

B. V. Kamath, K. Shivram and A. C. Shah 343 Determination of diclofenac sodium, famotidine and ketorolac tromethamine by flow injection analysis using dichloronitrophenol

E. Kiliç, F. Köseoğlu and M. A. Akay 347 The non-aqueous titrimetric assay of selected antibiotics using tetra-N-butylammonium hydroxide as titrant

D. J. Chapman, S. P. Joel and G. W. Aherne 353 Evaluation of a differential radioimmunoassay technique for the determination of morphine and morphine-6-glucuronide in human plasma

S. A. Wring, R. M. Rooney, C. P. Goddard, I. Waterhouse and W. N. Jenner 361 A sensitive radioimmunoassay, combined with solid-phase extraction, for the sub-nanogram per ml determination of ondansetron in human plasma

M. Berrabah, D. André, F. Prévot, A. M. Orecchioni and O. Lafont 373 GC–MS determination of phenobarbitone entrapped in poly-ε-caprolactone nanocapsules

M. J. Akhtar, S. Khan and M. Hafiz 379 High-performance liquid chromatographic assay for the determination of paracetamol, pseudoephedrine hydrochloride and triprolidine hydrochloride

S. Torrado, S. Torrado and R. Cadórniga 383 Comparison of assay methods by second-derivative spectroscopy, colorimetry and fluorescence spectroscopy of salicylic acid in aspirin preparations with a high-performance liquid chromatographic method

S. Miyairi, H. Shimada, N. Awata, J. Goto and T. Nambara 389 Novel immunoaffinity extraction for liquid chromatographic determination of major metabolities of 4-acetoxy-2-(4-methylphenyl) benzothiazole in plasma

T. Shintani, M. Takamoto, M. Sawada, H. Aishita and T. Nakagawa — 397 — Simultaneous determination of human neutrophil elastase inhibitor (ONO-5046) and its metabolite in plasma and urine by direct injection column-switching HPLC

S. Laganière and L. Goernert — 407 — Amrinone and *N*-acetylamrinone assay in human plasma using solid-phase extraction and reversed-phase chromatography

G. Kominami, H. Okabe, K. Imoda and K. Mizojiri — 413 — Combined high-performance liquid chromatography and radioimmunoassay for ceruletide and its metabolites in dog plasma and urine

Short Communications

M. J. Seddon, M. Spraul, I. D. Wilson, J. K. Nicholson and J. C. Lindon — 419 — Improvement in the characterization of minor drug metabolites from HPLC–NMR studies through the use of quantified maximum entropy processing of NMR spectra

A. Witek, P. Zawisza and L. Przyborowski — 425 — Determination of disopyramide in plasma by high-performance liquid chromatography

M. Koel and P. Nebinger — 429 — HPLC determination of serum ganciclovir using ultrafiltration, ultraviolet and fluorescence detection

Number 4

Review

H. A. H. Rongen, R. M. W. Hoetelmans, A. Bult and W. P. van Bennekom — 433 — Chemiluminescence and immunoassays

Full Papers

M. Moors, B. Steenssens, I. Tielemans and D. L. Massart — 463 — Solid-phase extraction of small drugs on apolar and ion-exchanging silica bonded phases: towards the development of a general strategy

T. F. Greber, T. V. Olah, J. D. Gilbert, A. G. Porras and M. Hichens — 483 — Determination of L-691, 121, a new Class III antiarrhythmic, and its principal metabolite in plasma by differential radioimmunoassay

K. C. Jindal, R. S. Chaudhary, A. K. Singla, S. S. Gangwal and S. Khanna — 493 — Dissolution test method for rifampicin–isoniazid fixed dose formulations

S. Levin and G. Tawil — 499 — Effect of surfactants on the diffusion coefficients of proteins, measured by analytical SPLITT fractionation (ASF) in the diffusion mode

M. Blanco, J. Coello, F. González, H. Iturriaga, S. Maspoch and X. Tomás — 509 — Spectrophotometric determination of pharmaceutical dosages by partial least-squares calibration

A. G. Barbone, B. Aparicio, D. W. Anderson, J. Natarajan and D. M. Ritchie — 515 — Reticulocyte measurements as a bioassay for erythropoietin

J. L. Burrows and K. V. Watson — 523 — Development and application of a calibration regression routine in conjunction with linear and non-linear chromatographic detector responses

H. Changqin, J. Shaohong and W. Kaimin — 533 — The chromatographic behaviour of cephalosporins in gel filtration chromatography, a novel method to separate high molecular weight impurities

L. Mälkki-Laine and A. P. Bruins — 543 — Structural characterization of the decomposition products of salbutamol by liquid chromatography–ionspray mass spectrometry

W. Naidong and J. W. Lee — 551 — Development and validation of a liquid chromatographic method for the quantitation of ibuprofen enantiomers in human plasma

N. Mano, Y. Oda, H. Ohe, N. Asakawa, Y. Yoshida and T. Sato — 557 — Resolution of 4-(4-chlorobenzyl)-2-(hexahydro-1-methyl-1H-azepin-4yl)-1 (2H)-phthalazinone enantiomers in plasma with frit-FAB LC–MS using a conalbumin column

Short Communications

W.-N. Wu and J. A. Masucci — 569 — Identification of two new suprofen metabolites in human urine

J. P. Zodda, J. A. Monteferrante and M. N. Eakins — 573 — Separation and simultaneous determination of methyl paraben and propyl paraben in Choletec® (kit for the preparation of Tc-99m mebrofenin) by high-performance liquid chromatography

I Announcements

Number 5

Review

N. W. Smith and M. B. Evans — 579 — Capillary zone electrophoresis in pharmaceutical and biomedical analysis

Full Papers

M. Spraul, J. K. Nicholson, M. J. Lynch and J. C. Lindon — 613 — Application of the one-dimensional TOCSY pulse sequence in 750 MHz ^{1}H-NMR spectroscopy for assignment of endogenous metabolite resonances in biofluids

M. C. Linhares and P. T. Kissinger — 619 — Pharmacokinetic studies using microdialysis probes in subcutaneous tissue: effects of the co-administration of ethanol and acetaminophen

J. A. Murillo and A. Alañón — 629 — New rapid assay for methicillin by spectrofluorimetry in pharmaceutical dosage forms

C. A. Georgiou, G. N. Valsami, P. E. Macheras and M. A. Koupparis — 635 — Automated flow-injection technique for use in dissolution studies of sustained-release formulations: application to iron(II) formulations

G. S. Clarke — 643 — The validation of analytical methods for drug substances and drug products in UK pharmaceutical laboratories

A. Parra, J. Garcia-Villanova, V. Rodenas and M. D. Gomez — 653 — First and second derivative spectrophotometric determination of cefoperazone and sulbactam in injections

D. Lessard, B. Comeau, A. Charlebois, L. Letarte and I. M. Davis — 659 — Quantification of GR90291 in human blood by high resolution gas chromatography–mass selective detection (HRGC–MSD)

E. I. A. Karim, K. E. E. Ibrahim, A. N. Abdelrahman and A. F. Fell — 667 — Photodegradation studies on chloroquine phosphate by high-performance liquid chromatography

P. D. Bryan and J. T. Stewart — 675 — Chromatographic analysis of selected tetracyclines from dosage forms and bulk drug substance using polymeric columns with acidic mobile phases

A. A. Nomeir, M. F. McComish, A. L. Mongan, J. M. Covey and N. F. Ferrala — 693 — Liquid chromatographic analysis, stability and protein binding studies of the anti-HIV agent benzoic acid, 2-chloro-5[[(1-methyl-ethoxy)thioxomethyl]amino]-,1-methylethyl ester

J. M. Lim, H. Kim, A. Marco, P. Mojaverian and C.-C. Lin — 699 — Liquid chromatographic determination of ceftibuten, a new oral cephalosporin, in human plasma and urine

T. V. Olah, J. D. Gilbert, A. Barrish, T. F. Greber and D. A. McLoughlin — 705 — A rapid and specific assay, based on liquid chromatography–atmospheric pressure chemical ionization mass spectrometry, for the determination of MK-434 (a 5α-reductase inhibitor) and its metabolites in plasma

Short Communications

G. W. Ponder and J. T. Stewart — 713 — Recovery of digoxin and related glycosides from an injection dosage form by liquid–liquid and solid-phase extraction

L. S. Wigman, H. Abdel-Kader and G. K. Menon — 719 — Size exclusion chromatography of poloxalene poloxamers: polyethylene glycol–polypropylene glycol co-polymers used to control cattle bloat

Number 6

Review

J. Šmisterová, K. Ensing and R. A. De Zeeuw — 723 — Methodological aspects of quantitative receptor assays

Full Papers

A. R. Lee and T. M. Hu — 747 — Determination of guaiphenesin in anti-tussive pharmaceutical preparations containing dextromethorphan by first- and second-derivative ultraviolet spectrophotometry

M. Heuermann and G. Blaschke — 753 — Simultaneous enantioselective determination and quantification of dimethindene and its metabolite *N*-demethyl-dimethindene in human urine using cyclodextrins as chiral additives in capillary electrophoresis

M. R. Euerby, C. M. Johnson, R. J. Lewis and S. C. Nichols — 761 — The S-oxidative degradation of a novel corticosteroid tipredane (INN). Part II—Detailed investigations into the primary S-oxidation of tipredane using achiral oxidants

L. A. McKown, W.-N. Wu and P. J. O'Neill — 771 — Characterization and identification of the metabolites of fenoctimine using *in vitro* drug metabolizing systems

K. Raghavan, A. Dwivedi, G. C. Campbell Jr, G. Nemeth and M. A. Hussain — 777 — A spectroscopic investigation of DuP 747 polymorphs

Z. Yu, Y. Xu and M. P. C. Ip — 787 — An ultra-sensitive electrochemical enzyme immunoassay for thyroid stimulating hormone in human serum

S. Chishima — 795 — A radioimmunoassay for TA-0910, a new metabolically stable thyrotrophin-releasing hormone analogue

L. Valvo, A. Paris, A. L. Savella, B. Gallinella and E. Ciranni Signoretti — 805 — General high-performance liquid chromatographic procedures for the rapid screening of natural and synthetic corticosteroids

J. Cummings, A. MacLellan, S. P. Langdon and J. F. Smyth — 811 — Stability and *in vitro* metabolism of the mitogenic neuropeptide antagonists [D-Arg^{1}, D-Phe^{5}, D-Trp^{7,9}, Leu^{11}]-substance P and [Arg^{6}, D-Trp^{7,9}, MePhe^{8}]-substance P (6–11) characterized by high-performance liquid chromatography

V. Brizzi and D. Corradini — 821 — Rapid analysis of somatostatin in pharmaceutical preparations by HPLC with a micropellicular reversed-phase column

G. A. Jacobson and G. M. Peterson — 825 — High-performance liquid chromatographic assay for the simultaneous determination of ipratropium bromide, fenoterol, salbutamol and terbutaline in nebulizer solution

M. Girard, K. Hannah, J. A. Menzies, G. Solomonraj and L. W. Whitehouse — 833 — Reversed-phase LC assay method for deoxycholate in influenza vaccine

K. Kobayashi, K. Chiba, M. Tani, Y. Kuroiwa and T. Ishizaki — 839 — Development and preliminary application of a high-performance liquid chromatographic assay for omeprazole metabolism in human liver microsomes

W. Naidong, A. M. Dzerk and J. W. Lee — 845 — Development and validation of an LC method for the quantitation of carbenicillin in human serum

Short Communication

K. M. Wasan, A. C. Hayman and G. Lopez-Berestein — 851 — Determination of dimyristoylphosphatidylglycerol in human serum by liquid–liquid extraction and reversed-phase liquid chromatography

Number 7

Full Papers

A. F. Casy and G. H. Dewar — 855 — Captopril and its probable contaminants: NMR and MS features of analytical value

J. Russell, J. Ness, M. Chopra, J. McMurray and W. E. Smith — 863 — The assessment of the HO · scavenging action of therapeutic agents

M. Salo, S. Sarna and H. Vuorela — 867 — Statistical evaluation of molecular descriptors and quantitative structure–property relationship studies of retinoids

J. A. Murillo, J. M. Lemus and L. F. García — 875 — Spectrofluorimetric analysis of cefuroxime in pharmaceutical dosage forms

Z. G. de Balugera, M. A. Goicolea and R. J. Barrio — 883 — Joint determination of todralazine and acetazolamide in human serum by differential pulse polarography

A. Kiss, J. Répási, Z. Salamon, Cs. Novák, G. Pokol and K. Tomor — 889 — Solid state investigation of mefloquine hydrochloride

I. Mahmood, S. H. Neau and W. D. Mason — 895 — An enzymatic assay for the MAO-B inhibitor selegiline in plasma

A. Van Overbeke, W. Baeyens, W. Van Den Bossche and C. Dewaele — 901 — Separation of 2-arylpropionic acids on a cellulose based chiral stationary phase by RP-HPLC

A. Van Overbeke, W. Baeyens, W. Van Den Bossche and C. Dewaele — 911 — Enantiomeric separation of amide derivatives of some 2-aryl-propionic acids by HPLC on a cellulose-based chiral stationary phase

V. Ulvi and H. Keski-Hynnilä — 917 — First-derivative UV spectrophotometric and high-performance liquid chromatographic analysis of some thiazide diuretics in the presence of their photodecomposition products

R. O. Epemolu, R. Ackerman, J. B. Porter, R. C. Hider, L. A. Damani and S. Singh — 923 — HPLC determination of 1,2-diethyl-3-hydroxypyridin-4-one (CP94), its iron complex [Fe(III) $(CP94)_3$] and glucuronide conjugate [CP94-GLUC] in serum and urine of thalassaemic patients

F. Benhamou-Batut, F. Demotes-Mainard, L. Labat, G. Vinçon and B. Bannwarth — 931 — Determination of flunitrazepam in plasma by liquid chromatography

H. P. J. M. Noteborn, F. J. Varossieau, M. Monshouwer, R. Brands and H. M. Pinedo — 937 — A reversed-phase high-performance liquid chromatographic method for the determination of Clanfenur in rat and human plasma

K. Gumbhir and W. D. Mason — 943 — Determination of *m*-hydroxymandelic acid, *m*-hydroxyphenyl-glycol and their conjugates in human plasma using liquid chromatography with electrochemical detection

Short Communications

A. Z. M. Wahbi, E. M. Hassan and A. A. Gazy — 951 — Use of isosbestic points of derivative absorption curves: the dissolution rate determination of Aspirin

F. A. Aly, F. Belal and M. I. Walash — 955 — Colorimetric determination of prenalterol hydrochloride in dosage forms

Number 8

Papers from the Eighth Annual American Association of Pharmaceutical Scientists Meeting, November 1993, Orlando, Florida, USA

959 — Editorial

A. A. Srigritsanapol and K. K. Chan — 961 — A rapid method for the separation and analysis of leaked and liposomal entrapped phosphoramide mustard in plasma

J.-S. Chiou and I. Ueda — 969 — Ethanol unfolds firefly luciferase while competitive inhibitors antagonize unfolding: DSC and FTIR analyses

I. Jääskeläinen and A. Urtti — 977 — Liquid chromatography determination of liposome components using a light-scattering evaporative detector

E. W. Tsai, S. D. Chamberlin, R. J. Forsyth, C. Bell, D. P. Ip and M. A. Brooks — 983 — Determination of bisphosphonate drugs in pharmaceutical dosage formulations by ion chromatography with indirect UV detection

K. P. Shah, J. Zhou, R. Lee, R. L. Schowen, R. Elsbernd, J. M. Ault, J. F. Stobaugh, M. Slavik and C. M. Riley — 993 — Automated analytical systems for drug development studies. I—A system for the determination of drug stability

General Papers

J. T. B. Strode III, L. T. Taylor, A. L. Howard, D. Ip and M. A. Brooks — 1003 — Analysis of felodipine by packed column supercritical fluid chromatography with electron capture and ultraviolet absorbance detection

E. M. Chellquist, R. Searle, D. L. Ladd and R. K. Hollister — 1015 — Binding constant determination of WIN 22169, a novel polymeric ligand

P. Amorusi, D. Lessard, S. K. Bansal, K. Selinger, A. Yacobi and A. P. Tonelli — 1023 — Analysis of enloplatin by liquid chromatography and of platinum by atomic absorption spectrometry in various biological fluids

Z. Shahrokh, G. Eberlein and Y. J. Wang — 1035 — Probing the conformation of protein (bFGF) precipitates by fluorescence spectroscopy

L.-S. Wu, C. Gerard, C. Koval, S. Rowe and M. A. Hussain — 1043 — Crystallinity, hygroscopicity and dissolution of moricizine hydrochloride hemihydrate

E. L. Hand, J. D. Gilbert, A. S. Yuan, T. V. Olah and M. Hichens — 1047 — Determination of MK-383, a non-peptide fibrinogen receptor antagonist, in human plasma and urine by radioimmunoassay

Short Communication

D. C. Rakestraw, D. A. Bilski and G. N. Lam — 1055 — Determination of linopirdine and its N-oxide metabolites in rat plasma by liquid chromatography

Number 9

Full Papers

R. M. E. Richards and D. K. L. Xing — 1063 — Determination of thymidine, uracil and *p*-aminobenzoic acid in bacteriological cultures by capillary zone electrophoresis

D. Boyd, J. R. B. Rodríguez, P. T. Blanco and M. R. Smyth — 1069 — Application of a Nafion-modified carbon paste electrode for the adsorptive stripping voltammetric determination of fenoterol in pharmaceutical preparations and biological fluids

R. G. Dickinson, A. R. King, M. A. Kelly, I. A. Kaltashov and C. Fenselau — 1075 — Excretion of 3-hydroxy-diflunisal as a monosulphate conjugate—identification using ESI-MS

V. Doyle and R. Devery — 1079 — A direct two-site peroxidase-based enzyme-linked immunosorbent assay for proinsulin

S. Luterotti and D. Maysinger — 1083 — Chemiluminometric determination of choline-related substances in pharmaceutical preparations by dot-blot

P. Saini, J. C. K. Lai and D. R. Lu — 1091 — FT–IR measurement of mercaptoundecahydrododecaborate in human plasma

M. C. Mahedero, F. Salinas and J. J. Aaron — 1097 — Determination of sulphanilamide in milk by first-derivative and second-derivative spectrofluorimetry

L. E. Bennett, W. N. Charman, D. B. Williams and S. A. Charman — 1103 — Analysis of bovine immunoglobulin G by capillary gel electrophoresis

T. Pérez-Ruiz, C. M. Lozano and V. Tomás — 1109 — Flow-injection determination of Novalgin using amperometric detection at a glassy carbon electrode

S. Levin, E. Klausner and S. Muchtar — 1115 — Characterization of submicron MCT o/w emulsions using sedimentation field-flow fractionation (FFF) with power field programming

A. Marsh and M. B. Evans — 1123 — Radioimmunoassay of drugs of abuse in hair. Part 1: Methadone in human hair, method adaptation and the evaluation of decontamination procedures

Z. H. Mohamed, S. M. Amer and A. M. El-Kousasy	1131	Colorimetric determination of cyclophosphamide and ifosphamide
Zhu Yongxin, C. Hendrix, R. Busson, G. Janssen, E. Roets and J. Hoogmartens	1137	Isolation and structural elucidation of an impurity of cefradine
D. Korakas, K. Valkó, C. Sakarellos, M. Sakarellos-Daisiotis and W. A. Gibbons	1141	Analytical and semi-preparative separation of diastereomeric lipidic amino acid conjugates
H. Wakabayashi, S. Yamato, M. Nakajima and K. Shimada	1147	Application of an electrochemical detector with a graphite electrode to liquid chromatographic determination of penicillamine and captopril in biological samples
M. J. Priston and G. J. Sewell	1153	Improved LC assay for the determination of mitozantrone in plasma: analytical considerations
C. M. Lai, B. Brogdon, C. Y. Quon and H. J. Pieniaszek Jr	1163	Determination of DuP 128, an ACAT inhibitor and its sulphoxide and sulphone metabolites in human plasma by liquid chromatography
A. R. Wafelman, M. C. P. Konings, H. Rosing, C. A. Hoefnagel, B. G. Taal, R. A. A. Maes and J. H. Beijnen	1173	High-performance liquid chromatographic determination of meta-iodobenzylguanidine in whole blood and plasma of cancer patients
G. S. Murkitt, E. Z. Trafankowska, M. Dunne and G. R. Manchee	1181	Fully automated assay for the determination of GR117289X in human plasma using mixed mode solid-phase extraction and high-performance liquid chromatography with ultraviolet detection
C. Pham-Huy, A. Sahui-Gnassi, V. Saada, J. P. Gramond, H. Galons, S. Ellouk-Achard, V. Levresse, D. Fompeydie and J. R. Claude	1189	Microassay of propranolol enantiomers and conjugates in human plasma and urine by high-performance liquid chromatography after chiral derivatization for pharmacokinetic study
Short Communications		
J. S. Millership	1199	Interference of mandelic acid with the determination of homatropine hydrobromide by second-order derivative spectroscopy
R. Boulieu and N. Bleyzac	1205	Stability of ganciclovir in blood samples
M. A. El-Ries, F. M. Abou Attia, F. M. Abdel-Gawad and S. M. Abu El-Wafa	1209	Preparation and characterization of Cu(II)–lignocaine complex. Indirect estimation of lignocaine in pharmaceutical preparations by atomic absorption spectroscopy (AAS)

Number 10

Full Papers		
M. Spraul, P. Neidig, U. Klauck, P. Kessler, E. Holmes, J. K. Nicholson, B. C. Sweatman, S. R. Salman, R. D. Farrant, E. Rahr, C. R. Beddell and J. C. Lindon	1215	Automatic reduction of NMR spectroscopic data for statistical and pattern recognition classification of samples
W. N. Moore and L. T. Taylor	1227	Analytical inverse supercritical fluid extraction of polar pharmaceutical compounds from cream and ointment matrices
P. C. Sontum and C. Christiansen	1233	Precision and accuracy of analysis of air-filled albumin microspheres using Coulter Multisizer Mark II
R. J. Forsyth and D. P. Ip	1243	Determination of imipenem and cilastatin sodium in Primaxin® by first order derivative ultraviolet spectrophotometry
P. C. Gunaratna, G. S. Wilson and M. Slavik	1249	Pharmacokinetic studies of α-difluoromethylornithine in rabbits using an enzyme-linked immunosorbent assay
K. Raghavan, G. Nemeth, D. Gray and M. A. Hussain	1259	Stabilization of an anthrapyrazole antitumour agent, DuP 937, on complexation with heptakis(2,6-di-O-methyl)-β-cyclodextrin in aqueous solution

T. Roy 1265 Fitting a straight line when both variables are subject to error: pharmaceutical applications

S. Scypinski and A. J. Ross 1271 Liquid chromatographic separation of zalcitabine and its stereoisomers

T. A. Walker 1277 A liquid chromatographic assay for urea in over-the-counter carbamide peroxide products

H. F. Deutsch, M. A. Evenson, P. Drescher, C. Sparwasser and P. O. Madsen 1283 Isolation and biological activity of aspidospermine and quebrachamine from an Aspidosperma tree source

J. Cummings, H. Fielding and W. R. Miller 1289 Pharmaceutical analysis of 8-chloroadenosine 3′, 5′-monophosphate

L. Šoltés, B. Sébille and P. Szalay 1295 Propafenone binding interaction with human α_1-acid glycoprotein: assessing experimental design and data evaluation

R. L. Tacey, W. J. Harman and L. L. Kelly 1303 Development of a highly sensitive and specific assay for plasma ethinylestradiol using combined extraction, liquid chromatography and radioimmunoassay

A. K. Seneviratne, A. L. Jayewardene and J. G. Gambertoglio 1311 Paired ion reversed-phase HPLC assay for the determination of iothalamic acid and para aminohippuric acid in urine

V. K. Dua, R. Sarin and V. P. Sharma 1317 Sulphadoxine concentrations in plasma, red blood cells and whole blood in healthy and *Plasmodium falciparum* malaria cases after treatment with Fansidar using high-performance liquid chromatography

Short Communications

D. L. Ross and C. M. Riley 1325 Dissociation and complexation of the fluoroquinolone antimicrobials—an update

P. Damiani, G. Ibáñez and A. C. Olivieri 1333 Zero-crossing first and second derivative synchronous fluorescence spectroscopic determination of aspirin metabolites in urine

Number 11

Analytical Survey

C. Hartmann, D. L. Massart and R. D. McDowall 1337 An analysis of the Washington Conference Report on bioanalytical method validation

Full Papers

J. M. Schuette, D. L. Cole and G. S. Srivatsa 1345 Development and validation of a method for routine base composition analysis of phosphorothioate oligonucleotides

E. Tarelli, P. H. Corran, B. R. Bingham, H. Mollison and R. Wait 1355 Lysine vasopressin undergoes rapid glycation in the presence of reducing sugars

R. Nellore and H. R. Bhagat 1363 Development and validation of capillary electrophoresis assay for Ribonuclease A

K. Takács-Novák, A. Avdeef, K. J. Box, B. Podányi and G. Szász 1369 Determination of protonation macro- and microconstants and octanol/water partition coefficient of the antiinflammatory drug niflumic acid

J. O. De Beer, C. V. Vandenbroucke and D. L. Massart 1379 Experimental design for the rapid selection of separation conditions for methyl and propyl parahydroxybenzoate, phenylephrine hydrochloride and chlorphenamine maleate by ion-pair liquid chromatography

Y. Sugawara and S. Nakamura 1397 Positive ion EI mass spectra of 2,3,5-trisubstituted-1,5-benzothiazepines, diltiazem, clentiazem and their fat-soluble metabolites

J. M. López, M. Redondo, T. Téllez and M. Morell 1411 A sensitive enzyme immunoassay for angiotensin II in serum

M. I. Walash, F. Belal, M. E. Metwally and M. M. Hefnawy 1417 A selective fluorimetric method for the determination of some 1,4-benzodiazepine drugs containing a hydroxyl group at C-3

L. W. Whitehouse, A. Menzies, B. Dawson, T. D. Cyr, A. W. By, D. B. Black and J. Zamecnik 1425 Mouse hepatic metabolites of ketoconazole: isolation and structure elucidation

J. Hempenius, G. Hendriks, J. Hingstman, C. K. Mensink, J. H. G. Jonkman and C. C. Lin 1443 An automated analytical method for the determination of felbamate in human plasma by robotic sample preparation and reversed-phase high performance liquid chromatography

M. J. McNulty, T. L. Page and D. L. Deal 1453 Development and validation of a chiral LC method for analysis of the four stereoisomers of 1045U85 in plasma

D. Fabre, F. Bressolle, J. M. Kinowski, O. Bouvet, F. Paganin and M. Galtier 1463 A reproducible, simple and sensitive HPLC assay for determination of ofloxacin in plasma and lung tissue. Application in pharmacokinetic studies

Short Communication

J. L. del Estal, M. L. Alvarez-Bujidos, R. Balaña Fouce, D. Ordóñez and J. G. Prieto 1471 The intestinal absorption of Luxabendazole in rats

Number 12

Full Papers

L.-G. Danielsson and Zhang Yu-Hui 1475 Mechanized determination of *n*-octanol/water partition constants using liquid–liquid segmented flow extraction

J. W. Muritu, I. O. Kibwage, C. K. Maitai and J. Hoogmartens 1483 Evaluation of tetracycline raw materials and finished products found on the Kenyan market

S. Y. Monté, I. Ismail, D. N. Mallett, C. Matthews and R. J. N. Tanner 1489 The minimal metabolism of inhaled 1,1,1,2-tetrafluoroethane to trifluoroacetic acid in man as determined by high sensitivity ^{19}F nuclear magnetic resonance spectroscopy of urine samples

M. Foradada, M. Gallardo and J. Estelrich 1495 Design and applications of a new fluorimetric assay of thioguanine in liposomes

K. Lohner, A. C. Sen, R. Prankerd, A. F. Esser and J. H. Perrin 1501 Effects of drug-binding of the thermal denaturation of human serum albumin

S. K. Branch, U. Holzgrabe, T. M. Jefferies, H. Mallwitz and M. W. Matchett 1507 Chiral discrimination of phenethylamines with β-cyclodextrin and *heptakis*(2,3-di-O-acetyl) β-cyclodextrin by capillary electrophoresis and NMR spectroscopy

K. P. Shah, Ming Chang and C. M. Riley 1519 Automated analytical systems for drug development studies. II—A system for dissolution testing

J. S. Grundy, R. Kherani and R. T. Foster 1529 Photostability determination of commercially available nifedipine oral dosage formulations

G. Egginger, W. Linder, G. Brunner and K. Stoschitzky 1537 Direct enantioselective determination of (R)- and (S)-propranolol in human plasma. Application to pharmacokinetic studies

H. Tsuchiya, S. Ohtani, K. Yamada, N. Takagi and T. Hayashi 1547 Determination of tetrahydro-β-carbolines in urine by high performance liquid chromatography with suppression of artefact formation

J. Y.-K. Hsieh, C. Lin, B. K. Matuszewski and M. R. Dobrinska 1555 Fully automated methods for the determination of hydrochlorothiazide in human plasma and urine

G. S. N. Lau and J. A. J. H. Critchley 1563 The estimation of paracetamol and its major metabolites in both plasma and urine by a single high-performance liquid chromatography assay

Short Communications

S. A. Wring, R. M. O'Neill, J. L. Williams, W. N. Jenner, M. J. Daniel, M. R. D. Gray, J. J. Newman, G. N. Wells and D. R. Sutherland — 1573 — The production and evaluation of a radioligand and antiserum for the radioimmunoassay of sub-nanogram per millilitre concentrations of lamivudine

M. S. García, C. Sánchez-Pedreño, M. I. Albero and V. Ródenas — 1585 — Determination of ampicillin or amoxycillin in pharmaceutical samples by flow injection analysis

B. J. Gurley, S. Zermatten and D. Skelton — 1591 — Determination of antipyrine in human serum by direct injection restricted access media liquid chromatography

C. Rodriguez, E. S. Estape-Wainwright, M. B. Sidhom, J. L. Cangiano and M. Martinez-Maldonado — 1597 — A simple spectrophotometric determination of submicromolar quantities of vanadium oxyions

A. K. Dash — 1601 — Determination of pseudoephedrine hydrochloride in dosage forms by liquid chromatography

I — Volume Contents, Author Index and Subject Index, Volume 12, 1994